## Updated Feb. 14,2020

As you are aware, an outbreak of a new straion confavirus, 2019-nCoV, presents an ongoing health threatacross the globe. We are sure you have already been watiligiengtly to respond to this threat. As information and ecommendations about 2019-nCoV continue to evolve, we owarn phasize our support of the CDC and other public hepatithners as the appropriate sourcie formation and guidance for health care organizations and providers.

The <u>CDC Infection Control webpater</u> health care professionals addresses several areas warthink worth emphasizing:

- Minimize the chance of exposurevolverking to identify patients before arr(inval), asking screening questions for patients callibrage pointments) and as early as possible apprimal (e.g., place signs for patients advisting to put on a mask if they have respirations).
- Implement triage procedures to defect2019-nCoVduring or before patient triage or registration (e.g., at the time of patient check-in) and ensure that all patients are asked abou presence of symptoms of a respiratory infection and history of travel to areas experiencing transmission of 2019-nCoV or contact proistable 2019-nCoV patients.
- If screening at triage is positive tamed patient becomes a person under investiga (PRUH), this should be communicated directly to tinecians who will care for the patient, prevention and control services, and other health care factivity according to a standard protocol.
- Adhere to standard contact and airborneecautions including eye protection.
- Monitor stock and the supply chairpersonal protective equipment.
- Manage visitor access and movement withher facility.
- Implement mechanisms and policies promptly alert key facility staff including infection control, health care epidemiology, facilitydership, occupational health, clinical laboratomyd frontline staff about known or suspected 2019 pactients.
- Identify specific staff to communicante collaborate with state or local public healthorities. Remember also that

This also is a good time to remind organizations of rourintections that are currently required the decrease the risk of transmission of any infectious agent. The set be in place regardless of an identified infectious threat:	nat will y new